

Semi-analytical study on the generic degeneracy for galaxy clustering measurements

Alejandro Guarnizo¹, Luca Amendola¹, Martin Kunz² and Adrian Vollmer¹

¹Institut Für Theoretische Physik, Ruprecht-Karls-Universität Heidelberg, Philosophenweg 16, 69120 Heidelberg, Germany

²Département de Physique Théorique and Center for Astroparticle Physics, Université de Genève, Quai E. Ansermet 24, CH-1211 Genève 4, Switzerland

Abstract. From the galaxy power spectrum in redshift space, we derive semi-analytical results on the generic degeneracy of galaxy clustering measurements. Defining the observables $\bar{A} = Gb\sigma_8$ and $\bar{R} = Gf\sigma_8$, (being G the growth function, b the bias, f the growth rate, and σ_8 the amplitude of the power spectrum), we perform a Fisher matrix analysis to forecast the expected precision of these quantities for a Euclid-like survey. Among the results we found that galaxy surveys have generically a slightly negative correlation between \bar{A} and \bar{R} , and they can always measure \bar{R} about 3.7 to 4.7 times better than \bar{A} .

Keywords. cosmology: observations, cosmology: theory, methods: statistical, surveys

1. Introduction

Future galaxy surveys will provide new opportunities to verify the current standard cosmological model, and also to constrain modified gravity theories, invoked to explain the present accelerated expansion of the universe. Before studying general parametrizations of dark energy, it is however important to understand first which quantities can be really observed. From this direction recently [Amendola et al. \(2013\)](#) shown that cosmological measurements can determine, in addition to the expansion rate $H(z)$, only three additional variables R , A and L , given by

$$A = Gb\delta_{m,0}, \quad R = Gf\delta_{m,0}, \quad L = \Omega_{m0}GY(1 + \eta)\delta_{m,0}. \quad (1.1)$$

with G is the growth function, b is the galaxy bias with respect to the dark matter density contrast, and $\delta_{m,0}$ is the dark matter density contrast today. The functions η (the anisotropic stress $\eta = -\frac{\Phi}{\Psi}$) and Y (the clustering of dark energy $Y \equiv -\frac{2k^2\Psi}{3\Omega_m\delta_m}$), describe the impact of the dark energy on cosmological perturbations. In [Amendola et al. \(2014\)](#), a Fisher analysis was made using galaxy clustering, weak lensing and supernovae probes, in order to find the expected accuracy with which an Euclid-like survey can measure the anisotropic stress η , in a model-independent way.

In this work we want to obtain some results on the intrinsic degeneracy on galaxy clustering measurements, using the quantities A and R . We use a flat Λ CDM fiducial model, with $\Omega_{m,0}h^2 = 0.134$, $\Omega_{b,0}h^2 = 0.022$, $n_s = 0.96$, $\tau = 0.085$, $h = 0.694$, $\Omega_k = 0$, Euclid-like survey specifications are used [Amendola et al. \(2013\)](#): we divided the redshift range $[0.5, 2.0]$ in 5 bins of width $\Delta z = 0.2$ and one of width $\Delta z = 0.4$; a spectroscopic error $\delta z = 0.001(1 + z)$, and a fraction of sky $f_{\text{sky}} = 0.375$; the bias b in the fiducial is assumed to be unity.

2. Fisher matrix for Galaxy Clustering

Observations of the growth rate f from large scale structures using Redshift Space Distortions (RSD), give a direct way to test different dark energy models, [Song & Percival \(2009\)](#), [Percival & White \(2009\)](#), [Racanelli *et al.* \(2013\)](#). Let us consider now the galaxy power spectrum in redshift space

$$P(k, \mu) = (A + R\mu^2)^2 = (\bar{A} + \bar{R}\mu^2)^2 \delta_{t,0}^2(k), \quad (2.1)$$

whit $\bar{A} = Gb\sigma_8$, $\bar{R} = Gf\sigma_8$, and we explicitly use $\delta_{m,0} = \sigma_8\delta_{t,0}$. The Fisher matrix is in general

$$F_{\alpha\beta} = \frac{1}{8\pi^2} \int_{-1}^1 d\mu \int_{k_{\min}}^{k_{\max}} k^2 V_{\text{eff}} D_{\alpha} D_{\beta} dk, \quad (2.2)$$

where $D_{\alpha} \equiv \frac{d \log P}{dp_{\alpha}}$, and V_{eff} is the effective volume of the survey

$$V_{\text{eff}} = \left(\frac{\bar{n}P(k, \mu)}{\bar{n}P(k, \mu) + 1} \right)^2 V_{\text{survey}}, \quad (2.3)$$

\bar{n} being the galaxy number density in each bin. We want to study the dependence on the angular integration in the Fisher matrix for the set of parameters $p_{\alpha} = \{A(z_{\alpha}), \bar{R}(z_{\alpha})\}$. The derivatives of the power spectrum are

$$D_{\alpha} = \frac{2}{\bar{A} + \bar{R}\mu^2} \{1, \mu^2\}. \quad (2.4)$$

We consider two cases depending on the behavior of V_{eff} , equation (2.3):

(a) “Enough data” $\bar{n}P(k, \mu) \gg 1$, then we have $V_{\text{eff}} \approx V_{\text{survey}}$ and the Fisher matrix could be written as

$$F_{\alpha\beta} \approx \frac{1}{2\pi^2} \int_{k_{\min}}^{k_{\max}} k^2 V_{\text{survey}} M_{\alpha\beta} dk, \quad (2.5)$$

where

$$M_{\alpha\beta} = 4 \begin{pmatrix} \frac{\sqrt{\bar{S}} + (\bar{A} + \bar{R}) \tan^{-1} \sqrt{P_1}}{\bar{A}^{3/2} \sqrt{\bar{R}}(\bar{A} + \bar{R})} & \frac{-\sqrt{\bar{S}} + (\bar{A} + \bar{R}) \tan^{-1} \sqrt{P_1}}{\bar{R}^{3/2} \sqrt{\bar{A}}(\bar{A} + \bar{R})} \\ \frac{-\sqrt{\bar{S}} + (\bar{A} + \bar{R}) \tan^{-1} \sqrt{P_1}}{\bar{R}^{3/2} \sqrt{\bar{A}}(\bar{A} + \bar{R})} & \frac{\bar{R}(3\bar{A} + 2\bar{R}) - 3(\bar{A} + \bar{R})\sqrt{\bar{S}} \tan^{-1} \sqrt{P_1}}{\bar{R}^3(\bar{A} + \bar{R})} \end{pmatrix} \quad (2.6)$$

being $\bar{S} = \bar{A}\bar{R}$ and $P_1 = \bar{R}/\bar{A}$.

(b) Shot-noise dominated $\bar{n}P(k, \mu) \ll 1$, then $V_{\text{eff}} \approx (\bar{n}P(k, \mu))^2 V_{\text{survey}}$ and since we are interested only in the μ dependence, we can write $V_{\text{eff}} \approx P(k, \mu)^2$. Then the Fisher matrix becomes

$$F_{\alpha\beta} \approx \frac{1}{2\pi^2} \int_{k_{\min}}^{k_{\max}} k^2 \delta_{t,0}^4(k) N_{\alpha\beta} dk, \quad (2.7)$$

with

$$N_{\alpha\beta} = 8 \begin{pmatrix} \bar{A}^2 + \frac{2\bar{A}\bar{R}}{3} + \frac{\bar{R}^2}{5} & \frac{\bar{A}^2}{3} + \frac{2\bar{A}\bar{R}}{5} + \frac{\bar{R}^2}{7} \\ \frac{\bar{A}^2}{3} + \frac{2\bar{A}\bar{R}}{5} + \frac{\bar{R}^2}{7} & \frac{\bar{A}^2}{5} + \frac{2\bar{A}\bar{R}}{7} + \frac{\bar{R}^2}{9} \end{pmatrix}. \quad (2.8)$$

We notice that in the two limiting cases above, we can move the matrices $M_{\alpha\beta}$ and $N_{\alpha\beta}$ outside of the integral, as for the fiducial model \bar{A} and \bar{R} do not depend on k . This




Figure 1. (color online). Confidence contours for \bar{A} and \bar{R} in the three cases: orange line V_{eff} , blue line $V_{\text{eff}} \approx V_{\text{survey}}$, and green line $V_{\text{eff}} \approx P(k, \mu)^2$.

means that, although the absolute size of the error ellipse depends on the integral, the relative size and orientation do not. In other words, we can obtain ‘generic expectations’ for the shape of the degeneracy between \bar{A} and \bar{R} from galaxy clustering surveys. These results are quite representative for the full range of \bar{A} and \bar{R} , i.e. galaxy surveys have

generically a slightly negative correlation between \bar{A} and \bar{R} , and they can always measure \bar{R} about 3.7 to 4.7 times better than \bar{A} , see Figure 1. In comparison to the results of Song & Kayo (2010), we remove the dependence on $\delta_{i,0}$, eq. (1.1), which is a quantity that depends on inflation or other primordial effects.

Acknowledgements

L.A., A.G. and A.V. acknowledge support from DFG through the project TRR33 “The Dark Universe”, A.G. also acknowledges support from DAAD through program “Forschungsstipendium für Doktoranden und Nachwuchswissenschaftler”. M.K. acknowledges financial support from the Swiss NSF.

References

- Amendola, L., Kunz, M., Motta, M., Saltas, I.D., & Sawicki, I. 2013, *Phys. Rev. D*, 87, 023501
 Amendola, L., Fogli, S., Guarnizo, A., Kunz, M., & Vollmer, A. 2014, *Phys. Rev. D*, 89, 063538
 Amendola, L., et. al. (Euclid Theory Working Group) 2013, *Living Rev. Relativity*, 16, 6
 Percival, W.J. & White, M. 2009, *MNRAS*, 393, 297-308
 Raccanelli, A., Bertacca, D., Pietrobon, D., Schmidt, F., Samushia, L., Bartolo, N., Dor, O., Matarrese, S., & Percival, W.J. 2013, *MNRAS*, 436, 89-100
 Song Y.-S., & Kayo I. 2009, *MNRAS*, 407, 1123-1127
 Song Y.-S., & Percival W. J. 2009, *J. Cosmol. Astropart. Phys.*, 0910, 004